Flora and Fauna Assessment

for the

Alteration and Additions

95 Bower Street, Manly





Document Information

PREPARED FOR:	ANNA WILLIAMS	
PREPARED BY:	Nicholas Skelton, B.Sc. (Hons), M.App.Sc. (Ecology and Vegetation Management) and Lucas Brown	
REVISION NO:	Final	
FILE NUMBER:	BS56TBR01	

Required Licences

NSW DEPARTMENT OF PRIMARY INDUSTRIES, ANIMAL RESEARCH AUTHORITY: 12/4838 OFFICE OF ENVIRONMENT AND HERITAGE, BIONET SENSITIVE SPECIES DATA LICENCE: CON97043 OFFICE OF ENVIRONMENT AND HERITAGE, BAM ASSESSOR: BAAS 1708

Approved for release by Director:

Nicholas Skelton, B.Sc. (Hons), M. App. Sc. GIS Environmental Consultants

Approval Date

6 May 2021

GIS Environmental Consultants

ABN: 66011504339 45 Austin Ave,

North Curl Curl, NSW 2099 Phone: (02) 9939 5129 Mobile: 0419 438 672

Email: ecology@ecology.net.au Web: <u>www.ecology.net.au</u>

Copyright GIS Environmental Consultants, All rights Reserved © 2021 GIS Environmental Consultants (Publisher) is the owner of the copyright subsisting in this publication. Other than as permitted by the Copyright Act and as outlined in the Terms of Engagement (for DA submission and assessment).

We acknowledge the traditional owners of this land the Gadigal people of the Eora nation and pay our respect to elders, past, present and emerging.

Table of Contents

Introduction4



6/5/21 Page 2 of 36

	1.1 Backg	round	2
	1.2 Legisla	ation Addressed by this Report	
	1.2.1	Environment Planning and Assessment Act 1979, EP&A Act	4
	1.2.2	Biodiversity Conservation Act 2016	4
	1.2.3	Biodiversity Conservation Act 2016, Threshold Test	
	1.2.4	Manly Local Environment Plan, MLEP 2013	(
	1.2.5	Manly Development Control Plan, DCP	
	1.2.6	State Environmental Planning Policies and Sydney Regional Environmental Plans	
	1.2.7	Federal Environment Protection and Biodiversity Conservation Act 1999, EPBC Act	
		al Definitions	
		nptions and Limitations	
		gered Bandicoot Population at North Head	
	1.5.1	Long-nosed Bandicoot (Perameles nasuta) Biology	
	1.5.2	Population Viability	
	1.5.3	Previous Years Survey Results for the Bushland Area of Habitat	
	1.5.4	Threats to the North Head Population	
		tudy Site	
		roposed Development	
	1.7.1	Plans and Documents Used	
2	Metho	ds	14
3	Findin	gs	1
Ū			
		nosed Bandicoot Use of Adjacent Land	
		g Bandicoot Habitat and Access	
	3.2.1	Existing Bandicoot Access to, and from, Adjacent Land	
	•	sed Bandicoot Habitat and Access	
	3.3.1	Proposed Bandicoot Access to, and from, Adjacent Land	
	3.3.2	During Construction Impacts to Bandicootsng Post-construction Impacts to Bandicoots	
	3.4.1		
	3.4.1 3.4.2	Artificial Lighting Noise	
	3.4.3	Avoiding Impacts	
	Trees 18	Avoiding impacts	1
		t and Presence of Other Flora and Fauna Species	18
	3.5.1	Non Threatened Fauna	
	3.5.2	Threatened Fauna	
	3.5.3	Threatened Plants	
	3.5.4	Non-Threatened Flora Species	
	3.6 Ecolog	gical Communities	
R /	•		21
IV	_	anges to Bandicoot Habitat and Access	
4	Photo	Page 1	22
5	Photo	Page 2	23
		G	
6	•	Assessment	
		f Significance (5-part test) for the Long-nosed Bandicoot Population	
	6.2 Manly	LEP 2013 Assessment of Clause 6.5 (3) & (4), Terrestrial Biodiversity	26
	6.2.1	Clause 6.5 (3) Assessment	
	6.2.2	Clause 6.5 (4) Assessment	
	6.3 State I	Environmental Planning Policies and Sydney Regional Environmental Plan	
	6.3.1	SEPP 19 Bushland in Urban Areas Assessment	
	6.3.2	SREP Sydney Harbour Catchment 2005 Assessment	
	6.3.3	SEPP Coastal Management 2018 Assessment	
	6.4 EPBC	Act 1999 Assessment	29
7	Conclu	usions	30
8		prative Conditions	
9	Manag	ement Recommendations	32



Pho	oto Page 3. Examples of Gaps in Boundary Fence	34
10	References and Relevant Literature	.35

1 Introduction

1.1 Background

This report identifies the ecological values and constraints at 95 Bower Street, Manly, then assesses the likely impact of a proposed development on the terrestrial flora, fauna and ecological communities, in particular, the Endangered Long-nosed Bandicoot population on North Head. The NSW Scientific Committee and OEH Profile has identified loss of habitat and change to habitat access as potential impacts and Key Threatening Processes to the Endangered Long-nosed Bandicoot population. This report also makes recommendations on ways to avoid or reduce impacts caused by the development.

1.2 Legislation Addressed by this Report

This section describes the Local, State and Federal legislation that provide the legal framework for approval of development and the protection and conservation of native flora and fauna that are relevant to this site.

1.2.1 Environment Planning and Assessment Act 1979, EP&A Act

The NSW Environment Planning and Assessment Act 1979 is the framework for approval of development in NSW. This proposal will be assessed under Part 4 of the EP&A Act, which requires the determining authority (usually Council) to not approve local development (Development Applications, DA's) without considering the heads of consideration in section 4.15, which requires the assessment of relevant legislation (Biodiversity Conservation Act, SEPP, LEPs, DCPs ect.)(4.15a), the environmental impact of the proposal (4.15b) and the suitability of the site for development (4.15c). Section 4 of this report addresses the BC Act and the relevant heads of consideration.

1.2.2 Biodiversity Conservation Act 2016

Section 7.2 of the Biodiversity Conservation Act requires that all local developments (Development Applications DAs, Part 4 EP&A Act):

- Implement the core purpose of the Act a hierarchy to "Avoid" and "Minimise" impacts; only then can "Offsets" be used for any residual impacts.
- Be assessed to determine whether they trigger the BOS Threshold Test specified in the Biodiversity Conservation Regulation 2017, which has two parts; the area of native vegetation that the proposal will impact and a check of whether the impact is within an area of mapped "biodiversity" on the Biodiversity values map; and
- Be assessed by a qualified ecologist to determine if there may be a positive a 5-Part Test of Significance as outlined in part 7.3 of the BC Act for each Threatened species or ecological community (listed in the schedules of the BC Act) or their habitats (listed in the schedules of the BC Act) that may occur on the site.
- Be assessed to determine if the proposal may impact on an Area of Outstanding Biodiversity Value (AOBV).

Developments that trigger the Threshold Test, or have a positive 5-Part Test of Significance, or impact on an AOBV, need to enter the Biodiversity Offset Scheme (BOS), and require that the Biodiversity Assessment Method (BAM) be applied and include a Biodiversity

GIS Environmental Consultants

6/5/21 Page 4 of 36

Development Assessment Report (BDAR) with the DA application. The proposal also needs to be assessed to determine if it may result in a Serious and Irreversible Impact (SAII).

If a development application does not meet the threshold or any other triggers, then a smaller report is still required to address the "heads of consideration" (section 4.15 of the EP&A Act), 5-part Test of Significance as required by the Manly DCP, SEPPs and Local Council's LEP/DCP requirements.


1.2.3 Biodiversity Conservation Act 2016, Threshold Test

This proposal is **not** considered to meet the BC Act threshold as;

- 1. The lot size is less than 1ha and there is less than 0.25ha of native vegetation being removed. **and**
- 2. The proposal will not directly or indirectly impact a declared Area or Outstanding Biodiversity Significance (AOBV) or an area mapped as having high biodiversity value on the "Biodiversity Values Map". See figure 1 below. **and**
- 3. There is not likely to be a significant affect (5-part test of significance test in Section 7.3, BC Act) on any Threatened species or ecological community or their habitat, as has been determined by section 6.1 of this report.

Therefore, the proposal does not need a Biodiversity Development Assessment Report (BDAR).

The Threshold Test, 5-Part Test, assessment of AOBV, heads of consideration, SEPP LEP/DCP requirements are all assessed in section 6 of this report.



Ref: https://www.lmbc.nsw.gov.au/Maps/index.html?viewer=BOSETMap



6/5/21 Page 5 of 36

1.2.4 Manly Local Environment Plan, MLEP 2013

The Manly (Northern Beaches Council) Local Environment Plan's (2013) 'Terrestrial Biodiversity Map' identifies this property as having high terrestrial biodiversity value. Development applications in the mapped area require consideration of Clause 6.5 (3) and (4) 'Terrestrial Biodiversity' (Manly LEP 2013). Development proposals need to be consistent with the objectives of this Clause and include appropriate measures to avoid, minimise or mitigate possible impacts of the development on biodiversity.

Extract from MLEP 2013

6.5 Terrestrial biodiversity

- (1) The objective of this clause is to maintain terrestrial biodiversity by:
 - (a) protecting native fauna and flora, and
 - (b) protecting the ecological processes necessary for their continued existence, and
 - (c) encouraging the conservation and recovery of native fauna and flora and their habitats.
- (2) This clause applies to land identified as "Biodiversity" on the Terrestrial Biodiversity Map.
- (3) Before determining a development application for development on land to which this clause applies, the consent authority must consider:
 - (a) whether the development is likely to have:
 - (i) any adverse impact on the condition, ecological value and significance of the fauna and flora on the land, and
 - (ii) any adverse impact on the importance of the vegetation on the land to the habitat and survival of native fauna, and
 - (iii) any potential to fragment, disturb or diminish the biodiversity structure, function and composition of the land, and
 - (iv) any adverse impact on the habitat elements providing connectivity on the land, and
 - (b) any appropriate measures proposed to avoid, minimise or mitigate the impacts of the development.
- (4) Development consent must not be granted to development on land to which this clause applies unless the consent authority is satisfied that:
 - (a) the development is designed, sited and will be managed to avoid any significant adverse environmental impact, or
 - (b) if that impact cannot be reasonably avoided by adopting feasible alternatives—the development is designed, sited and will be managed to minimise that impact, or
 - (c) if that impact cannot be minimised the development will be managed to mitigate that impact.

These sections of the Manly LEP are addressed within this report.

1.2.5 Manly Development Control Plan, DCP

Manly Development Control Plan 2013 Section 2.1.15 'Threatened Flora and Fauna Assessment of Significance Report, including the Long-nosed Bandicoot, and Little Penguins' requires the assessment of the significance of impact on threatened species, populations or ecological communities or their habitats.

Manly DCP 2013

DCP extract 2.1.15 'Threatened Flora and Fauna Assessment of Significance Report'

Objective 1) To ensure the assessment of any significant effect on threatened species, populations or ecological communities or their habitats (as listed in the Threatened Species Conservation Act 1995) in accordance with Section 5A of the Environmental Planning and Assessment Act 1979 (now superseded by section 7.3 of the Biodiversity Conservation Act 2016).

DCP extract 5.4.2 'Threatened Species and Critical Habitat'

Any development of land with known habitat for threatened species must consider the likely impacts of the development and whether further assessment needs to be undertaken by a Species Impact Statement.

DCP 3.3.1 Landscape Character



6/5/21 Page 6 of 36

- a) The design, quantity and quality of open space should respond to the character of the area. In particular:
- iv) In areas of habitat for the long-nosed bandicoot: (see paragraph 5.4.2), landscape design must include native plant species to provide new and/or improved low dense clumping habitat to provide for potential foraging and nesting. The planting schedule should comprise species such as Lomandra sp. Dianella sp., Banksia spinulosa, Caustis sp., Xanthorrhoea sp., Isolepis sp., Juncus sp., Adiantum sp., Calochlaena sp., Callistemon sp., Grevillea juniperina, Gleichenia sp., Grevillea 'Robyn Gordon' and tussocky native grasses (eg. Kangaroo Grass)

DCP Extract - Schedule 1 - Map D - Areas where Assessment of Significance is required (for Little Penguins and/or Long Nosed Bandicoots)



This report includes a Test of Significance (5-part test) for the Endangered Long-nosed Bandicoot population at North Head and other Threatened Species, Populations or Endangered Ecological Communities that may be impacted by the proposal. The Impact Assessment section of this report addresses these clauses.

1.2.6 State Environmental Planning Policies and Sydney Regional Environmental Plans

The SEPPs and SREPs which are relevant to Northern Beaches LGA and which may be relevant to this proposal are SEPP 19 Bushland in Urban Areas and Sydney Regional Environmental Plan (Sydney Harbour Catchment) (SREP SHC) 2005 (which amends SEPP No 56 Sydney Harbour Foreshores and Tributaries).

State Environmental Planning Policy No 19—Bushland in Urban Areas

State Environmental Planning Policy 19 – Bushland in Urban Areas (SEPP 19) is an NSW government policy that aims to protect and preserve bushland within urban areas. The policy applies to naturally vegetated land adjacent to Council reserves.

Sydney Regional Environmental Plan (Sydney Harbour Catchment) 2005

The plan aims to establish a balance between promoting a prosperous working harbour, maintaining a healthy and sustainable waterway environment and promoting recreational access to the foreshore and waterways. It establishes planning principles and controls and

GIS Environmental Consultants

6/5/21 Page 7 of 36

consolidates and replaces the following instruments: - Sydney Regional Environmental Plan No. 22 - Parramatta River (SREP 22); - Sydney Regional Environmental Plan No. 23 - Sydney and Middle Harbour (SREP 23); and amends State Environmental Planning Policy No. 56 Sydney Harbour Foreshores and Tributaries (SEPP 56).

The area to the south of the heavy black boundary line on the figure below is within the Foreshores and Waterways Area and includes the whole of North Head, St Patrick's Estate, Manly Boatshed and Manly Wharf. See image below. The whole of Manly Cove is zoned as W2 Environment Protection Zone. Wetlands are mapped along the majority of Manly Coves' foreshore.




Image: Strategic Foreshores and Waterways Area - Part of Sheet 4 SREP SHC

The ecological Aim of the SREP is to ensure the protection, maintenance and rehabilitation of watercourses, wetlands, riparian lands, remnant vegetation and ecological connectivity.

The Impact Assessment section of this report assesses the specific ecological matters that

are to be considered.

Coastal Management SEPP 2018

The new SEPP Coastal Management 2018 combines SEPP 14 (Coastal Wetlands), SEPP 26 (Littoral Rainforests) and SEPP 71 (Coastal Protection) and clause 5.5 of the Standard Instrument into one integrated policy. These policies have been repealed. This SEPP defines four coastal management areas and specifies the assessment of development within these management areas.

The Coastal Management SEPP also maps Coastal Wetlands and Littoral Rainforest (EEC) and areas within proximity of Coastal Wetland and Littoral Rainforest and includes controls for development within those areas.

1.2.7 Federal Environment Protection and Biodiversity Conservation Act 1999, EPBC Act

There is currently no memorandum of understanding agreement between the State and Federal government regarding the need to apply the EPBC Act 1999.

The Commonwealth Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) will need detailed assessment if the proposal is considered likely to have an impact on a

GIS Environmental Consultants

6/5/21 Page 8 of 36

'Matter of National Environmental Significance (MNES), thus providing a trigger for referral of the proposal to the Department of the Environment and Water Resources. Matters of national environmental significance identified in the Act are; world heritage properties; national heritage places; RAMSAR wetlands; nationally threatened species and communities; migratory species protected under international agreements; the Commonwealth marine environment; and nuclear actions.

Section 4 of this report addresses this requirement.

1.3 General Definitions

5-Part Test of Significance (5-Part Test) - Assessment under Section 7.3 of the BC Act to determine whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats. The minister has provided a guide under 7.3(2) titled Threatened Species Assessment Guidelines.

BC Act - NSW Biodiversity Conservation Act 2016 contains the lists of threatened species, the definitions of the threatened ecological communities, the 5-part Test of Significance and the Biodiversity Offset Scheme (BOS). There are associated Biodiversity Conservation regulations which refers to the BAM.

Clearing – clearing of native vegetation including; cutting down, felling, uprooting, thinning or otherwise removing native vegetation, killing destroying, poisoning, ringbarking or burning <u>native vegetation</u> and includes and includes establishment and maintenance of bushfire protection Asset Protection Zones (APZ) inner and outer zones.

Direct Impacts - are impacts that directly affect habitat, ecosystems and individuals. They include, but are not limited to, death, trampling, poisoning of the animal/plant itself and the removal of vegetation and suitable habitat. When applying each factor, consideration must be given to all of the likely direct impacts of the proposed activity or development during construction. As defined by the 2018 Threatened Species Assessment Guidelines.

DPI – NSW government of Department of Primary Industries

EPA Act (EP&A Act) – NSW Environment Planning and Assessment Act 1979, controls development in NSW, includes the requirement to consider SEPPs, LEPs, DCPs, BC Act 2016.

EPBC Act – Federal Environment Protection and Biodiversity Conservation Act 1999.

Indirect Impacts - occur when project-related activities affect species, populations or ecological communities in a manner other than direct loss. Indirect impacts can include loss of individuals through starvation, exposure, predation by domestic and/or feral animals, loss of breeding opportunities, loss of shade/shelter, deleterious hydrological changes, increased soil salinity, erosion, inhibition of nitrogen fixation, weed invasion, fertiliser drift, or increased human activity within or directly adjacent to sensitive habitat areas. Indirect impacts may occur after construction during the life of the development, e.g. escape of garden plants, excess nutrients and changes in fire frequency and grazing. As with direct impacts, consideration must be given, to all of the likely indirect impacts of the proposed activity or development (2006 DECC Assessment of Significance Guidelines)

LEP – Local Environment Plan, a local planning instrument for each Council area.

Native Vegetation - is defined in the LLS Act as any plants native to NSW including trees, understory plants or groundcover plants including wetland. Marine vegetation is protected by the Fisheries Act.

OEH – NSW Office of Environment and Heritage, formerly NPWS, DEC, DECC and DECCW. Government organisation responsible for the conservation of native flora and fauna.

Property – The lot(s) that are the subject of the proposal. In this report, this is the same as the Study Area, the Subject Site and "site".

6/5/21 Page 9 of 36



Proposal – The works/actions that are proposed on the property that is the subject of the development application.

Protected Fauna - refers to any native bird, mammal, reptile or frog in NSW.

Site - In this report this is the same as the Study Area and the Subject Site and the property.

Study Area - means the subject site and any additional areas which are likely to be affected by the proposal, either directly or indirectly. The study area should extend as far as is necessary to take all potential impacts into account. In this report, this is the same as the Subject Site, the property and "site".

Subject Site - means the area directly affected by the proposal. In this report, this is the same as the Study Area, the property and "site".

Threatened Species - refers to those species listed in the schedules of the Biodiversity Conservation Act 2016 as "Critically Endangered", "Endangered" or "Vulnerable".

For definitions that are relevant to the Assessment of Significant test see the Appendices.

1.4 Assumptions and Limitations

- This document only assesses the impacts of the proposal described in this report and shown on Map 1 and the cited plans.
- This report does not take into account the cumulative impact of other developments on this property or on adjacent land.
- This report does not include assessment of soil suitability or European/Aboriginal heritage.
- It can never be proven that other Threatened Species have not, do not or will not use the site as habitat. The conclusions drawn in this report are a result of testing, observation and experience.
- This report describes the habitat and species of the site at the time of the field survey.
 Vegetation, habitat and legislation will change over time and therefore the findings of this report are only relevant for 6 months.
- This report should be read in its entirety and no part should be taken out of context.
- No responsibility is accepted for use of any part of this report in any other context or for any other purpose or by third parties.

1.5 Endangered Bandicoot Population at North Head

The main species of interest on this site is the Long-nosed Bandicoot, *Perameles nasuta*, (Geoffrey 1804) and in particular, the Endangered population at North Head, Manly, which is known to occur in the vicinity of the Subject Site.

The Final Determination (TSC Act Scientific Committee 1997) for the listing of this population in the schedules of the Threatened Species Conservation Act described the population as:



- "P. nasuta was once widespread in the Sydney region but many formerly recorded populations have become extinct. The North Head population is now isolated and disjunct."
- "...the North Head population of P. nasuta is in immediate danger of extinction."
- "...the North Head P. nasuta population is of significant conservation value on the grounds that it is:
 - A disjunct population

GIS Environmental Consultants

6/5/21 Page 10 of 36

- One of the few surviving populations within the Sydney region
- A population which has been the subject of a number of scientific studies, and is thus an important reference population
- Accorded considerable value by the local community, and thus serves to promote conservation more generally"

The Office of Environment and Heritage has identified 25 priority actions to help recover the Long-nosed Bandicoot population on North Head in New South Wales (as of July 2013). These priority actions relate to OEH, Northern Beaches Council and other determining authorities developing, implementing and continuing the fox, feral cat and rabbit control program, weed control program, monitoring program, community awareness program, collecting mortality data, finalising and reviewing Long-nosed Bandicoot Recovery Plan, and Sydney Harbour National Park Fire Management Strategy and Plan of Management.

1.5.1 Long-nosed Bandicoot (Perameles nasuta) Biology

Description: The Long-nosed Bandicoot (*Perameles nasuta*) is a solitary nocturnal marsupial that grows to a size of between 850 and 1100 g, 310 to 425 mm in head and body length, and with a tail length of 120 to 155 mm (Stoddart 1995). The males are larger than females. These bandicoots characteristically dark, greyish-brown above and creamy white below. The forefeet and upper surfaces of the hind feet are also creamy white (NPWS 2000b). The muzzle is long and pointed and the ears are markedly larger and more pointed than short-nosed bandicoots of the genus Isoodon, such as the other bandicoot that lives in Sydney, the Southern Brown Bandicoot (Stoddart 1995).

Distribution: Long-nosed Bandicoots are locally common along the east coast of Australia and adjacent mountains from north-eastern Queensland to south-western Victoria. This Endangered population is restricted to the relatively isolated area of habitat on North Head in the Manly Local Government Area, south of Addison Road (NSW Scientific Committee 2003). See above the DCP extract, Schedule 1 - Map D - Areas where Assessment (test) of Significance is required. There is another Threatened population in the inner western part of Sydney.

Habitat: At North Head, Long-nosed Bandicoots inhabit, to varying extents, all of the habitat types available including woodlands, scrub, heath open areas and the urban landscape. Recent research indicates that urban areas are important for the population and that there are individuals who live their entire lives within the urban area. Long-nosed Bandicoots prefer sites with sandy soils, as well as with low undergrowth and leaf litter cover and does not have a particular preference for proportion of canopy cover (Chambers & Dickman 2002). This species depends on a mosaic of vegetation types at a landscape level, including feeding grounds in patches of moist, soft soil located close to shelter with an abundance of invertebrates (Scott et al. 1999). These types of habitat can be found in both bushland and urban environments including native vegetation and residential gardens. Resting and nesting habitat is low, dense vegetation or litter where a bandicoot can take shelter in during the day. Bandicoot diggings are more abundant in areas of moist, soft soils close to cover.

Individuals build diurnal nests that are typically made in a shallow hole or depression on ground surface and are lined with leaf litter and dry grasses. The entrance to each nest is closed when occupied making them generally difficult to locate. Nests on North Head+ have been found in a variety of habitat types, such as at the base of large trees and within tall grasses including residential backyards (Scott 1995; Scott *et al.* 1999). Long-nosed Bandicoots typically have more than 1 nest that is in regular use within their territory (Chambers & Dickman 2002). It is expected that bandicoots in the wild may live up to 2 to 2.5 years.

6/5/21 Page 11 of 36



Home Range Size: Home range size of an individual Long-nosed Bandicoots have been recorded at 1.3ha (+-0.2 S.E. 50% KDE) for females (n=5) and 1.1ha (50% KDE) for a male. Animals tended to maintain exclusive and relatively stable core home ranges, although overlap of non-homes ranges was common (Hope 2012).

Diet: Long-nosed Bandicoots feed on invertebrates, plants, tubers, fungi and vertebrates (Menkhorst & Knight 2004, Scott et al. 1999, Claridge 1993). Invertebrates mostly include insects from the orders Coleoptera and Hymenoptera (> 80%). Plants preferred are mainly the leaves and stems of monocotyledons (>76%). Fungi are consumed in a high proportion (> 63%), mostly those hypogeal from the family Zygomicetes, in particular the species *Glomus fueglanum*. Vertebrates contribute little to bandicoots' diet, but include skinks, birds and sometimes eggs of the Eastern Water Dragon (Scott et al. 1999).

Breeding: At North Head, Long-nosed Bandicoots were recorded breeding from June to March (Scott 1995), however mating can occur throughout the year. The average recorded litter size for the North Head population is 2.3 babies (Stoddart 1995). In productive years, females may have up to 4 litters.

Litters are typically 2-3 (>76%), with the young weaned at about 7 weeks and reaching maturity at 20 weeks. Females tend to overlap their home ranges (i.e., 1.7 ha) throughout the year, as well as to reduce their size during the breeding season. By contrast, home ranges of males (i.e., 4.4 ha) only overlap during the breeding season, as they also enlarge their home ranges (Scott et al. 1999, Menkhorst & Knight 2004).

1.5.2 Population Viability

There have been many studies on this population over the last 20 years including; microchipping, radio tracking, extensive trapping, diet analysis, population viability estimation (Banks, 2000; Banks, 2004; Chambers and Dickman, 2002; Hughes and Banks, 2006; Hughes and Banks, 2010; Lenehan and Banks, 2004; Scott, Hume, and Dickman, 1999). There is ongoing biannual monitoring program by the Office of Environment and Heritage (OEH; formerly DECCW, DECC and NPWS) in collaboration with Manly Council and Sydney University.

Every two years there is a more extensive Long-nosed Bandicoot trapping survey conducted in the bushland part of North Head. This survey does not include the urban environment part of North Head, which is now known to have bandicoot residence. These areas are likely to be the same population. Population viability estimates within the urban environment most recently occurring in November 2012 and March 2013. This urban study utilised 14 transects across Eastern Hill and used the same methods as the current study by NPWS within the bushland habitat on North Head (Hughes and Banks, 2010).

In May 2015, a total of 152 individual Long-nosed Bandicoots were trapped at North Head, compared to 99 in 2014, and 71 in 2010. The sex ratio of the bushland population is relatively even. Under current conditions, the North Head Long-nosed Bandicoot population has a 62% chance of persisting after 50 years. This figure has dropped from 80% in since the previous PVA, due to the slightly higher sex specific adult mortality rates used in the current PVA's (Price & Banks, 2015). The latest PVA analysis determined that the population is stable and has been for the last few years. It has been calculated that only a small loss to the population could cause the local population to become extinct.

1.5.3 Previous Years Survey Results for the Bushland Area of Habitat

See below text results from the Long-Nosed Bandicoot Urban Monitoring Program.

• A total of 34 (14 males and 19 females, adults 72%) individual Long-nosed Bandicoots were trapped in the urban area of Manly, in May 2016, compared with 31 (19 males and 12 females, 75% adult) in March 2013.

6/5/21 Page 12 of 36



- Four of the females captured had 1-2 young in their pouch.
- 25% of the total population on North Head are living within the urban environment with 28 – 45 individuals in the urban environment compared with 120 - 140 individuals in the bushland environment (NPWS).
- There are individuals with their home range within the urban environment
- Individuals were trapped across Eastern Hill and down to Ashburner Street.

1.5.4 Threats to the North Head Population

The major threats to this population are thought to be vehicle traffic, loss of habitat through development and, to a lesser degree, predation by dogs, cats and foxes. Other threats include inbreeding depression, loss of genetic variation, the risk of catastrophic events (such as bushfires or disease), inappropriate fire regimes, clearing of native vegetation and invasion of native plant communities by Bitou Bush. Bandicoots are also susceptible to infection by cats carrying the disease toxoplasmosis. If urban developments keep reducing the area of accessible habitat available it is likely to result in population decrease and the likelihood of the population becoming extinct in the near future (i.e., 20 years) range between 31% and 46%.

The removal of habitat or prevention of access to habitat on a site may constitute a significant impact to the conservation of the threatened population and may require modifications to the development so there is no significant impact or a more extensive assessment in the form of a Species Impact Statement or modification of the proposal.



6/5/21 Page 13 of 36

1.6 The Study Site

The Site is the whole of Lot 81 D.P 8076, known as 95 Bower St Manly. The Study Site is a corner lot with an area of 799.7m².

The property contains an existing ground and first floor and several lawns that front the property. See Map 1 for plans showing the existing site, habitat and access.

There are areas of urban habitat surrounding the site, and extensive areas of bushland reserve on North Head, mostly Sydney Harbour National Park and Sydney Harbour Federation Trust land to the south-east of the property. Bower Streets runs along the northern border and residential properties border the site to the south and west. The eastern boundary of the property is adjacent to college street. Further to the north is Cabbage Tree Bay Aquatic Reserve. Approximately 50m away is St Pat's estate, which contains good quality habitat and bushland.

The geographic co-ordinates of the site are 33.801° S and 151.293° E.

1.7 The Proposed Development

This report addresses a DA for:

- The alterations and additions to the dwelling
- Landscaping

For further information on locations, extent of the development and details of the proposal, see Map 1.

The plans and documents used for this report are:

1.7.1 Plans and Documents Used

Title	Author	Re v	DWG./Doc. No./Ref. No.	Date Modified or Accessed
Survey Plan	Waterview Surveying Services	Α	1196detail1	18/09/20
Site Plans	Your Beautiful Home	В	SK05	16/03/21
Landscape Master Plan	Harrison's Landscaping	Α		27/04/21

2 Methods

The Site was inspected on the 17th of March 2021 by experienced and qualified ecologist Nicholas Skelton, Lucas Brown and Joshua Drane for a total of 3-person hours. Nicholas Skelton has 20 years of experience in Flora and Fauna surveys in the Sydney Metropolitan area and has completed over 200 bandicoot surveys and assessments in Manly for NPWS, Council, SHFT and private landowners. The field survey searched for evidence of all Threatened Species, Populations and Endangered Ecological Communities that are known to, or that may have potential habitat within the site, especially the Endangered population of Long-nosed Bandicoot.

Existing and potential foraging, resting, and nesting Long-nosed Bandicoot habitat was determined and quantified and is shown on Map 1. Existing bandicoot access to, from and within the site was also identified and mapped.

6/5/21 Page 14 of 36



The plans referenced within this report were assessed to determine the amount and type of habitat and the access that would be altered as a result of the proposal. Map 1 shows the change in the amount of habitat and access.

The habitat potential of the site for bandicoots was determined by detailed onsite assessment of the access, shelter and food sources. The recent use of the property by bandicoots was determined by an ecologist with extensive experience in bandicoot survey in urban environments, by searching for diggings, scats, frequently used trails and boundaries were thoroughly searched for accessibility by bandicoots. The road reserve and accessible parts of nearby properties were searched for evidence of bandicoot activity and habitat value. Photographs were taken of the site. The findings from other reports from nearby surveys and studies were also used to provide additional habitat use information. Habitat for other Threatened species was searched for.

3 Findings

3.1 Long-nosed Bandicoot Use of Adjacent Land

During the field survey evidence was found of Bandicoots using the site and adjacent path of public access. It is likely that bandicoots also use the adjacent properties and nearby bushland areas for resting and breeding as well as foraging due to the good quality habitat. Bandicoots have been recorded regularly in the locality. These animals are all from the Endangered Longnosed Bandicoot population at North Head.

3.2 Existing Bandicoot Habitat and Access

The site currently contains 355 m² of bandicoot foraging habitat (44% of the site) in the lawn and garden areas at the front and rear of the site. No resting or nesting bandicoots were observed on the property during the site survey.

Existing access is provided from the north west, at various intervals along a fence which borders the property, and otherwise blocks bandicoot access. Access is also provided from gaps under the eastmost gate, and the driveway gate on the northern side of the property. Further access is provided by the fence bordering the east of the property. Gaps in the fence bordering the south of the property provides access to the retaining wall from adjacent properties, lot 83 and 84. The fence that borders the Western lawn to the south provides access from adjacent property, lot 82. Bandicoots can currently access all the habitat on the property.

There is a low retaining wall that borders the front of the poperty, mostly blocking access except for under the eastern most gate and the driveway gate. There is a high retaining wall that borders the rear of the property, which bandicoots may access from the eastern side of the property and through gaps in the wall. They will only be able to access the appropriate habitat this way by running along the wall towards the eastern most lawn.

Map 1 shows the existing bandicoot habitat and access at the site. Areas shown on green are areas of bandicoot foraging habitat.

3.2.1 Existing Bandicoot Access to, and from, Adjacent Land

See Dark Blue arrows on Map 1 that show bandicoot access and Red lines show barriers bandicoot access.

GIS Environmental Consultants

6/5/21 Page 15 of 36

Adjacent Land	Bandicoot Access and Habitat		
North– Bower St.	Good quality habitat – Road reserve, footpath and Council road reserve. Access is good through the north of the property.		
South – Neighbouring property	Good quality habitat – Access through the southern fence.		
East – College St.	Good quality habitat – Road reserve, footpath and council reserve to the south east. Access from the east of the property.		
West – Neighbouring property	Good quality habitat – Footpath and council reserve. Access from the western fence.		

3.3 Proposed Bandicoot Habitat and Access

Map 1 summarises the existing and proposed bandicoot habitat and access to, from and within the site. Areas shown on green are areas of bandicoot foraging habitat.

To avoid and minimise impact to bandicoot habitat we have had a meeting with the architchcts the owner and the landscape designer, where we recommended appropriate acess points and habitat areas.

The proposal will result in a loss of 77 m² of foraging habitat from with the property. There will also be temporary (during construction) removal of most of the habitat as a result of demolition, excavation, storage of materials and stockpiling.

The proposal will provide 278 m² (35% of the site) of potential foraging habitat in the form of landscaped areas in the front, rear and sides of the property.

There will be access every 2m along the fence that borders College Street, the fence that borders the property to the south, from the start of the eastern lawn, three points of access from the western fence and one point of access from the northern gate.

The proposal will change access for bandicoots. The following proposals have been made to maintain provide additional access (see Magenta arrows and Light Blue lines on Map 1).

- Access is recommended from Bower Street through a gap under the new reconfigured front gate. Any new front gates are to have a minimum 150mm gap under the full length of the base;
- Access to the rear yards of the property is to be maintained along the eastern side of the proposed dwelling.
- Any gates along the through access path from the front to rear are to have 150mm gaps for bandicoots;
- Where there is no change in soil levels, bandicoot access is to be maintained through the existing gaps in the fences along the northern and southern boundaries;
- Any replacement boundary fences are to have 150mm-300mm gaps every two metres along the fence.

3.3.1 Proposed Bandicoot Access to, and from, Adjacent Land

See Dark Blue arrows on Map 1 that show proposed bandicoot access and Red lines that show bandicoot access barriers.

GIS Environmental Consultants

6/5/21 Page 16 of 36

Adjacent Land	Bandicoot Access and Habitat	
North– Bower St.	Good quality habitat – Road reserve, footpath and council reserve to the south east. Access from the east and north of the property retained, however with less points of access form the north.	
South – Neighbouring property	Good quality habitat – However, 29 and 31 Reddall street are no longer accessible due to the height of the proposed southern wall. Access to 35 Reddall street is retained.	
East – College St.	Good quality habitat – Road reserve, footpath and council reserve. Access retained from the east of the property.	
West – Neighbouring property	Good quality habitat – Footpath and council reserve. Access to be retained the from western fence.	

3.3.2 During Construction Impacts to Bandicoots

Likely impacts during construction are:

- Temporary restriction of access to habitat for bandicoots and other fauna due to skips, site fences and storage of materials. **Medium impact**;
- Temporary potential hazards to bandicoots and other fauna including falling into open pits and drowning hazards. Low risk, this is not very likely at this site;
- Temporary additional traffic movement around the street. Low risk.

The proposal will temporarily remove a large portion of the habitat on the site during construction due to demolition, potential excavation and material storage.

Measures to ameliorate these potential impacts are discussed in the Recommendations and Ameliorative Conditions sections of this report.

3.4 Ongoing Post-construction Impacts to Bandicoots

3.4.1 Artificial Lighting

Artificial lights may deter bandicoots from using areas of habitat. Multiple lights, bright lights or motion sensing lights are likely to be detrimental to bandicoots. Impacts can include increase in fox, dog, owls and cats predation

3.4.2 Noise

Noise is likely to deter bandicoots. There is currently very little information regarding the amount or type of noise that has impact. Potential noise sources are alarms, people and music.

3.4.3 Avoiding Impacts

No amplified music, reduced lighting, careful design of light shades to reduced light spill, use of multiple low level LEDs rather than overhead lighting, use of warmer red end spectrum lighting use of light coloured surfaces and flooring.

6/5/21 Page 17 of 36



Trees

3.5 Habitat and Presence of Other Flora and Fauna Species

3.5.1 Non Threatened Fauna

Species	Common name
Physignathus lesueurii	Eastern Water Dragon
Trichoglossus moluccanus	Rainbow Lorikeet
Lampropholis guichenoti	Garden Skink
Ocyphaps lophotes	Crested Pigeon
Manorina melanocephala	Noisy Miner
Trichosurus vulpecula	Brush-tailed possum
Alectura lathami	Brush Turkey

3.5.2 Threatened Fauna

Threatened Grey-headed Flying-foxes regularly feed from the fig and micro-bats regularly fly over this property. There are OEH BioNet records of these species occurring in the locality. There is a large amount foraging habitat in the locality for these species. No evidence of any flying-foxes or microbats roosting at the site.

Approximately ten years ago, the local population of Eastern Pygmy Possum and Brown Antechinus became extinct, and the local population of native Bush Rat population became non-viable or locally extinct. The Sydney Harbour Federation Trust in collaboration with Sydney University, then reintroduced the Eastern Pygmy Possum, Brown Antechinus and the native Bush Rat to North Head. The three species are all breeding, and the local populations are becoming established, with the Bush Rat population being the most successful. So far as they are out competing the introduced Black Rat and now the population covers most of North Head.

The native Bush Rat can possibly be found in the urban area; however, it is unlikely that the Antechinus or the Pygmy Possum occur in the urban area yet.

The native Bush Rat, Antechinus and Eastern Pygmy-possum have been recorded along Darley Road, adjacent to the north of St Patricks

3.5.3 Threatened Plants

There are local populations of the endangered Magenta Lillypilly, Syzygium paniculatum and Sunshine Wattle (Acacia terminalis subsp. terminalis) on North Head and the Magenta Lillypilly has been recorded in the nearby Bower Street Gully Reserve but it does not occur on this site. No evidence was found of any other Threatened Species, Population or Endangered Ecological Community on this property at the time of the survey. None of the other six (6) endangered and twenty-eight (28) vulnerable fauna species that occur in the Manly area where found on the site or have important habitat on the site.

6/5/21 Page 18 of 36



3.5.4 Non-Threatened Flora Species

There are not enough native species on the site or the correct structure for the vegetation on the site to represent any native vegetation community see Table 1 below for a list of native plants found at the site. No Endangered Ecological Community occurs on the site.

Table 1. Plant Species in Northern Wildlife Corridor

Genus and Species	Family	Habit	Order	Common Name	Status
				Creeping	
Commelina cyanea	COMMELINACEAE	Herb	MONOCOTYLEDON	Christian	Local native
Cyathea cooperi	CYATHEACEAE	Fern	FERN	Straw Tree Fern	Local native
Ficus rubiginosa	MORACEAE	Tree	DICOTYLEDON	Port Jackson Fig	Local native
Glochidion ferdinandi var. ferdinandi	EUPHORBIACEAE	Tree	DICOTYLEDON	Cheese Tree	Local native
Pittosporum				Sweet	
undulatum	PITTOSPORACEAE	Tree	DICOTYLEDON	Pittosporum	Local native
Syzygium australe	MYRTACEAE	Tree	DICOTYLEDON	Bush Cherry	Local native
Brugmansia					
suaveolens	SOLANACEAE	Shrub	DICOTYLEDON	AngelsTrumpet	Planted
Callistemon Hybrid	MYRTACEAE	Shrub	DICOTYLEDON	Bottle Brush	Planted
Hydrangea					
macrophylla	SAXIFRAGACEAE	Herb	DICOTYLEDON	Hydrangea	Planted
Nerium oleander	APOCYNACEAE	Shrub	DICOTYLEDON	Oleander	Planted
				Canary Island	
Phoenix canariensis	ARECACEAE	Palm	MONOCOTYLEDON	Date Palm	Planted
	1050105:5		MONOCOTI (1 = 2 2) :	Dwarf Date	.
Phoenix roebelenii	ARECACEAE	Palm	MONOCOTYLEDON	Palm	Planted
Acetosa sagittata	POLYGONACEAE	Vine	DICOTYLEDON	Turkey Rhubarb	Weed
Ageratina adenophora	ASTERACEAE	Herb	DICOTYLEDON	Crofton Weed	Weed
Asparagus officinalis	ASPARAGACEAE	Herb	MONOCOTYLEDON	Asparagus	Weed
Cerastium glomeratum	CARYOPHYLLACEAE	Herb	DICOTYLEDON	Mouse Ear Chick Weed	Weed
Chlorphytum					
comosum	LILLIACEAE	Herb	MONOCOTYLEDON	Spider Plant	Weed
Ehrharta erecta	POACEAE	Grass	MONOCOTYLEDON	Ehrharta	Weed
Ipomea indica	CONVOLVULACEAE	Vine	DICOTYLEDON	Morning glory	Weed
Lantana camara	VERBENACEAE	Shrub	DICOTYLEDON	Lantana	Weed
Ligustrum sinense	OLEACEAE	Shrub	DICOTYLEDON	Privet - narrow leaved	Weed
	0.4.001.001.001.00	\ , <i>r</i>	DIOCENII EE CO	Japanese	
Lonicera japonica	CAPRIFOLIACEAE	Vine	DICOTYLEDON	Honeysuckle	Weed
Macfadyena unguis-	DICNONIACEAE	Vina	DICOTYLEDON	Cat's Claw	Wood
cati Morus alba	BIGNONIACEAE MORACEAE	Vine Tree	DICOTYLEDON DICOTYLEDON	Creeper	Weed Weed
			FERN	White Mulberry	
Nephrolepis cordifolia Nothoscordum gracile	DAVALLIACEAE ALLIACEAE	Fern	MONOCOTYLEDON	Fishbone Fern Onion Weed	Weed Weed
Notifoscordum gracile	ALLIACEAE	Herb	IVIONOCOTYLEDON		vveeu
Ochna serrulata	OCHNACEAE	Shrub	DICOTYLEDON	Ochna, Mickey Mouse Plant	Weed
Olea europaea subsp.					
africana	OLEACEAE	Shrub	DICOTYLEDON	African Olive	Weed
B. J. C. J. J. J. J.	LIDTIOAGEAE		DIOOTVI EDON	Asthma Weed,	VA /
Parietaria judaica	URTICACEAE	Herb	DICOTYLEDON	Pellitory	Weed
Senna pendula	CAESALPINIOIDEAE	Shrub	DICOTYLEDON	Cassia	Weed
Stenotaphrum				5 # 1 6	
secundatum	POACEAE	Grass	MONOCOTYLEDON	Buffalo Grass	Weed
Syngonium sp.	ARACEAE	Herb	MONOCOTYLEDON	Syngonium	Weed

Some of the weed species within this property require the land owner to take action as describe in the Greater Sydney Regional Weed Management Plan 2017-2022 https://greatersydney.lls.nsw.gov.au/ data/assets/pdf file/0010/722368/Greater-Sydney-Regional-Weed-Mgmt-Plan-FINAL-Sept-2019.pdf.

Table 2. Priority Weeds from Greater Sydney Regional Weed Management Plan



6/5/21 Page 19 of 36

Species	Action Requred
Asparagus aethiopicus	Person must not move, import into the State or sell.
Lantana camara	Person must not move, import into the State or sell.

3.6 Ecological Communities

The NSW Threatened Species Conservation Act (TSC), 1995 and the Federal Environment Protection and Biodiversity Conservation (EPBC) Act 1999, both list Threatened Ecological Communities. Threatened ecological communities can be either Vulnerable (VEC) or Endangered (EEC) or Critically Endangered (CEEC) Ecological Communities under the TSC Act. The Federal Act lists only Endangered or Critically Endangered Ecological Communities. These communities are likely to become extinct in nature unless the circumstances and factors threatening their survival cease to operate. The listing is most commonly referred to as a determination, which is a several page definition of the community written by a scientific committee and listed in the schedules of the Act.

During the site survey, the likelihood of Endangered Ecological Communities occurring on the site was determined using a three-step approach: 1. Has the community been recorded in the locality? 2. Is there a sufficient density of characteristic species on the site? 3. Does the environmental description in the Determination fit the site?

Littoral Rainforest Endangered Ecological Community

Littoral rainforest in the NSW North Coast, Sydney Basin and South East Corner Bioregions endangered Ecological Community (EEC) is generally a closed forest, the structure and composition of which is strongly influenced by proximity to the ocean. The plant species in this ecological community contain some rainforest species with evergreen mesic or coriaceous leaves, however the site does not contain enough species to be considered Littoral Rainforest. Planted Littoral Rainforest is likely to occur in the nearby Bower Street Gully Reserve, approximately 500m away.



6/5/21 Page 20 of 36

Map 1: Changes to Bandicoot Habitat and Access






Figure 1. Back of the house looking to western boundary, showing the retaining wall and stairs to be removed at far end



Figure 2. Existing bandicoot access along the southern boundary of the property

6 Impact Assessment

This Test of Significance is in accordance with the Threatened Species Assessment Guidelines recommended for use by Manly Development Control Plan 2013 (MDCP 2013) in section 2.1.15.2 (a).

6.1 Test of Significance (5-part test) for the Long-nosed Bandicoot Population

Part 7.3 of the BC Act, Test of Significance (5-part test) for impact of the proposed development on the Long-nosed Bandicoot Population at North Head

(a) in the case of a threatened species, whether the proposed development or activity is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,

Response:

The Long-nosed Bandicoot population on North Head is listed in Schedule 1, Part 2, Division 4 of the BC Act 2016 as an Endangered Species Population.

The local population is viable at least in the short term. In May 2015, a total of 152 individual Long-nosed Bandicoots were trapped at North Head, compared to 99 in 2014, and 71 in 2010. The sex ratio of the bushland population is relatively even. Under current conditions, the North Head Long-nosed Bandicoot population has a 62% chance of persisting after 50 years. This figure has dropped from 80% in since the previous PVA, due to the slightly higher sex specific adult mortality rates used in the current PVA's (Price & Banks, 2015). The latest PVA analysis determined that the population is stable and has been for the last few years.

The site currently contains 355 m² of bandicoot foraging habitat (44% of the site) in the lawn and garden areas at the front and rear of the site. The proposal will result in a loss of 77 m² of foraging habitat from with the property. The proposal will provide 278 m² (35% of the site) of potential foraging habitat in the form of landscaped areas in the front, rear and sides of the property (see before and after Green areas on Map 1).

This population is restricted to North Head, which has an area of 385 ha, of which only a negligible amount will be gained due to the proposed development. The change in bandicoot habitat is of a scale that is not likely to lead to the significant reduction in the population size or reproduction success of individuals, the population or their habitat. The proposal is not likely to have a significant negative effect on the life cycle of this population such that the viability of the population is compromised and placed at risk of extinction.

- (b) in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:
 - (i) is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or
 - (ii) is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,

Response:

The Long-nosed Bandicoot Population at North Head is listed as a threatened population and not an Endangered or Critically Endangered Ecological Community; therefore, this question is not applicable.

6/5/21 Page 24 of 36



- (c) in relation to the habitat of a threatened species, population, or ecological community:
 - i) the **extent** to which habitat is likely to be removed or modified as a result of the action proposed, and

Response:

The site currently contains 355 m² of bandicoot foraging habitat (44% of the site) in the lawn and garden areas at the front and rear of the site. The proposal will result in a loss of 77 m² of foraging habitat from with the property. The proposal will provide 278 m² (35% of the site) of potential foraging habitat in the form of landscaped areas in the front, rear and sides of the property (see before and after Green areas on Map 1).

This population is restricted to North Head, which has an area of 385 ha, of which only a negligible amount will be gained due to the proposed development. The change in the extent of bandicoot habitat is of a scale that is not likely to lead to the significant reduction in the population size or reproduction success of individuals, the population or their habitat. The proposal is not likely to have a significant negative effect on the life cycle of this population such that the viability of the population is compromised and placed at risk of extinction.

ii) whether an area of habitat is likely to become **fragmented or isolated** from other areas of habitat as a result of the action proposed, and

Response:

Access to habitat on the property will be decreased by the proposal. See Map 1 for proposed bandicoot habitat and access to, from and within the site. Existing access to habitat surrounding the site will not be changed as a result of the proposal.

iii) the **importance** of the habitat to be removed, modified, fragmented or isolated to the **long-term survival** of the species, population or ecological community in the locality

Response:

This population is restricted to North Head, which has an area of 385 ha, of which only a negligible amount will be gained due to the proposed development.

The change in bandicoot habitat is of a scale that is not likely to lead to the significant reduction in the population size or reproduction success of individuals, the population or their habitat. The proposal is not likely to have a significant negative effect on the life cycle of this population such that the viability of the population is compromised and placed at risk of extinction.

(d) whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),

Response:

There is no Area of Outstanding Biodiversity on the site. The proposal will not directly or indirectly effect any Area of Outstanding Biodiversity Value.

(e) whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.

Response:

The following Key Threatening Processes are relevant to the Site and/or the proposal

6/5/21 Page 25 of 36



Clearing of native vegetation: The proposal will remove 2 local native Cheese trees. This is a small impact. Therefore, the proposal is not likely to increase this or any other listed Key Threatening Process.

Conclusion to the 5-Part Test of Significance on the Endangered populations of Longnosed Bandicoots.

It is not likely that the proposal will have a significant impact on the Endangered Long-nosed Bandicoot population at North Head. Entry into the Biodiversity Offset Scheme (BOS) and further assessment in the form of a Biodiversity Development Assessment Report (BDAR) is not considered necessary for this population.

6.2 Manly LEP 2013 Assessment of Clause 6.5 (3) & (4), Terrestrial Biodiversity

Manly LEP 2013 'Terrestrial Biodiversity Map' shows the subject property is located within an area identified as of 'Terrestrial Biodiversity'.

Therefore Clause 6.5 of MLEP 2013 applies to this Development Application and the objectives of the clause and in particular points (3) and (4) must be considered in regard to this proposal.

6.2.1 Clause 6.5 (3) Assessment

- a) Whether the development is likely to have:
 - i. Any adverse impact on the condition, ecological value and significance of the fauna and flora on the land?

Response: It is likely that bandicoots use the adjacent properties and nearby bushland areas for foraging and probably resting and breeding. Long-nosed Bandicoots have been recorded regularly in the locality.

Other fauna that are likely to use site include possums, Rainbow Lorikeets, Noisy Minors, Sulphur Crested Cockatoo, Garden Skinks, Eastern Water Dragon and Brush turkeys, Crested Pigeon, Grey-headed Flying-fox, Eastern Water Dragons and a Diamond Python.

The site currently contains 355 m² of bandicoot foraging habitat (44% of the site) in the lawn and garden areas at the front and rear of the site. The proposal will result in a loss of 77 m² of foraging habitat from with the property. The proposal will provide 278 m² (35% of the site) of potential foraging habitat in the form of landscaped areas in the front, rear and sides of the property (see before and after Green areas on Map 1).

Access to the habitat will not change for birds or arboreal mammals such as possums that are using the site.

Access to the habitat for bandicoots (See Dark Blue arrows on Map 1) will be decreased by the proposal. Proposals have been made to maintain some access from the existing site.

No evidence was found of any other Threatened Species, Populations or Endangered Ecological Communities utilising this property.

Based on the information gathered and the assessments of potential impacts of the proposal on flora and fauna in section 3 of this report, it is considered that the proposal will not have any adverse impact on the condition, ecological value and significance of the fauna and flora on the land.

ii. Any adverse impact on the importance of the vegetation on the land to the habitat and survival of native fauna?

Response: The vegetation on the site is medium quality habitat for a wide range of fauna including reptiles, birds and mammals. Brush-tailed and Ring-tailed Possums are likely to occur.

GIS Environmental Consultants

6/5/21 Page 26 of 36

There is no native vegetation community on the property. The proposal will remove some of the existing lawn area in the front of the property.

Based on the findings and assessment of the impact of this proposal on flora and fauna in this report, fauna habitat is not likely to be adversely impacted by the proposal and the habitat is not likely to be important habitat for these species due to the higher quality bushland habitat to the east in Sydney Harbour National Park. The proposed development will not have any adverse impact on the importance of the vegetation on the land to the habitat and survival of native fauna.

It is recommended that local native species be used in landscaping on the property to improve that habitat value of the vegetation to native fauna.

iii. Any potential to fragment, disturb or diminish the biodiversity structure, function and composition of the land?

Response: Based on the findings and assessment of the impact of this proposal on flora and fauna in sections 3 and 4 of this report, the proposed development will not significantly fragment, disturb or diminish the current biodiversity structure, function and composition of the subject site. See section 4 for details.

iv. Any adverse impact on the habitat elements providing connectivity on the land?

Response: The site is in a residential area surrounded by other private properties with a discontiguous canopy of trees. Access to the habitat will not change for birds or arboreal mammals such as possums that are using the site. Access to the habitat for bandicoots (See Dark Blue arrows on Map 1) will not be changed by the proposal. Recommendations have been made to provide additional access (see Magenta arrows on Map 1).

Based on the findings and assessment of the impact of this proposal on flora and fauna in of this report, this proposal will not adversely impact on the habitat elements providing connectivity to other areas of suitable habitat.

b) Are there appropriate measures proposed to avoid, minimise or mitigate the impacts of the development?

Response: This report makes recommendations for appropriate measures to avoid, minimise or mitigate the impacts of the development. See the Ameliorative Conditions and Management Recommendations sections of this report for further information.

6.2.2 Clause 6.5 (4) Assessment

a) Is the development designed, sited and will be managed to avoid any significant adverse environmental impact? OR

Response: The development has been designed to utilise the site topography and surrounding landscape. The proposal will retain most existing access for bandicoots. The current and proposed access to the site is shown on Map 1. The recommendations and ameliorative conditions in this report provide measures to manage and mitigate impacts.

b) If the impact cannot be reasonably avoided by adopting feasible alternatives—is the development designed, sited and will be managed to minimise that impact? OR

Response: The development has been designed to utilise the site topography and surrounding landscape. The proposal will retain access along all boundaries of the property for movement of bandicoot species. The recommendations and ameliorative conditions in this report provide measures to manage and mitigate impacts.

c) If that impact cannot be minimised—will the development will be managed to mitigate that impact?

GIS Environmental Consultants

6/5/21 Page 27 of 36

Response: N/A

Conclusion to the Assessment of Clause 6.5 of the MLEP

The proposal is consistent with the requirements of Clause 6.5 of the MLEP 2013 and is not considered to have a significant adverse impact on terrestrial biodiversity

6.3 State Environmental Planning Policies and Sydney Regional Environmental Plan

6.3.1 SEPP 19 Bushland in Urban Areas Assessment

The site does not contain natural vegetation with intact structure and floristics and therefore does not fit the definition of Bushland as described in SEPP 19 (Bushland in Urban Areas 1986).

The proposed works, with the amelioration recommendations described in this report, will have a very low impact on the environment, they will not disrupt any fauna corridor, they will not endanger plant or animal species, they will not cause significant erosion and they will not change the accessibility or recreational value of bushland. The proposed works therefore are considered generally to meet the objectives of SEPP 19.

- a) to protect the remnants of plant communities which were once characteristic of land now within an urban area,
- b) to retain bushland in parcels of a size and configuration which will enable the existing plant and animal communities to survive in the long term,
- c) to protect rare and endangered flora and fauna species,
- d) to protect habitats for native flora and fauna,
- e) to protect wildlife corridors and vegetation links with other nearby bushland,
- f) to protect bushland as a natural stabiliser of the soil surface,
- g) to protect bushland for its scenic values, and to retain the unique visual identity of the landscape,
- h) to protect significant geological features,
- i) to protect existing landforms, such as natural drainage lines, watercourses and foreshores,
- j) to protect archaeological relics,
- k) to protect the recreational potential of bushland,
- I) to protect the educational potential of bushland,
- m) to maintain bushland in locations which are readily accessible to the community, and
- n) to promote the management of bushland in a manner which protects and enhances the quality of the bushland and facilitates public enjoyment of the bushland compatible with its conservation.

6.3.2 SREP Sydney Harbour Catchment 2005 Assessment

The site is not included within the Sydney Harbour Catchment Map and therefore assessment with respect to SREP Sydney Harbour Catchment is not required. However, this proposal is consistent with the matters to consider in respect to the biodiversity, ecology and environment protection of the site.

6/5/21 Page 28 of 36



6.3.3 SEPP Coastal Management 2018 Assessment

The site is mapped as Costal Environment Area in the Coastal Management SEPP 2018. The site is not mapped as containing Littoral Rainforest, Coastal Wetland, Proximity to Littoral Rainforest or Proximity to Coastal Wetland.

- (a) The proposed development should have a neutral or beneficial effect on the quality of water entering the waterways
- (b) The proposed development should protect and enhance terrestrial and aquatic species, populations and ecological communities and, in particular, should avoid physical damage and shading of aquatic vegetation (such as seagrass, saltmarsh and algal and mangrove communities)
- (c) The proposed development should promote ecological connectivity between neighbouring areas of aquatic vegetation (such as seagrass, saltmarsh and algal and mangrove communities)
- (d) The proposed development should avoid indirect impacts on aquatic vegetation (such as changes to flow, current and wave action and changes to water quality) as a result of increased access
- (e) The proposed development should protect and reinstate natural intertidal foreshore areas, natural landforms and native vegetation
- (f) The proposed development should retain, rehabilitate and restore riparian land
- (g) The proposed development on land adjoining wetlands should maintain and enhance the ecological integrity of the wetlands and, where possible, should provide a vegetative buffer to protect the wetlands
- (h) The cumulative environmental impact of development
- (i) Whether sediments in the waterway adjacent to the development are contaminated, and what means will minimise their disturbance.
- (a) to protect the natural and cultural values of waters in this zone,
- (b) to prevent damage or the possibility of longer term detrimental impacts to the natural and cultural values of waters in this zone and adjoining foreshores,
- (c) to give preference to enhancing and rehabilitating the natural and cultural values of waters in this zone and adjoining foreshores,
- (d) to provide for the long-term management of the natural and cultural values of waters in this zone and adjoining foreshores.

6.4 EPBC Act 1999 Assessment

Flora, fauna and ecological communities within Manly, which are listed under the EPBC Act: Eastern Suburbs Banksia Scrub is listed as Endangered.

Grey-headed Flying-fox (*Pteropus poliocephalus*) is listed as Vulnerable.

Littoral Rainforest and Coastal Vine Thickets of Eastern Australia is listed as Critically Endangered.

Sunshine Wattle (Acacia terminalis subsp. terminalis) is listed as Endangered.

Seaforth Mintbush (*Prostanthera marifolia*) is listed as Critically Endangered.

Pimelea curviflora var. curviflora is listed as Vulnerable.

North Head is listed as a National Heritage Place. About 277ha, at Manly, comprising the whole of the headland, to Low Water, south of a line commencing at Low Water

Page 29 of 36



north of Collins Beach on the alignment of the north-west boundary of Lot 2763 DP752038, then easterly via that alignment and boundary and then following the north-westerly boundaries of Lot 2774 DP752038 Lot 2728 DP752038, Lot 2764 DP752038 and Lot 2763 DP752038 to the most northerly point of Lot 2763 DP752038, then generally easterly via the north-east and northern boundaries of Lot 2763 DP752038 and the alignment of the latter segment to Low Water. Excluded is the North Head Sewage Treatment Plant being the whole of Lot 1 DP604428.

The only matters of relevance to this proposal are migratory species, threatened species and communities and national heritage places. North Head is on the National Heritage List as of 12 May 2006; however, the listing does not include this part of North Head. This Endangered Bandicoot Population is not listed in this Act. The relevant matters of National Environmental Significance have been considered. This proposal is not considered likely to have an impact on any matter of National Environmental Significance and referral is not required.

7 Conclusions

The site survey found evidence of the Long-nosed Bandicoot using the front and rear garden at the site. It is likely that bandicoots use this and adjacent properties and nearby bushland areas for foraging and probably resting and breeding. Long-nosed Bandicoot have been recorded regularly in the locality.

Other fauna that are likely to use the site include possums, Rainbow Lorikeets, Noisy Minors, Garden Skinks and Brush Turkeys, Crested Pigeon, Grey-headed Flying-fox and Eastern Water Dragons.

The site currently contains 355 m² of bandicoot foraging habitat (44% of the site) in the lawn and garden areas at the front and rear of the site. The proposal will result in a loss of 77 m² of foraging habitat from with the property. The proposal will provide 278 m² (35% of the site) of potential foraging habitat in the form of landscaped areas in the front, rear and sides of the property (see before and after Green areas on Map 1).

Access to the habitat will not change for birds or arboreal mammals such as possums that are using the site.

Access to the habitat for bandicoots (See Dark Blue arrows on Map 1) will be decreased by the proposal. Proposals have been made to maintain some access from the existing site.

No evidence was found of any other Threatened Species, Populations or Endangered Ecological Communities utilising this property.

The development is unlikely to have a significant impact on the conservation of any Endangered Population, Threatened Species or Endangered Ecological Community. The proposal does not meet the BC Act Threshold Test. Further assessment of the impact of this proposal in the form of a Biodiversity Development Assessment Report (BDAR) not recommended in relation to this development application at this site. The ecological impact is not considered an unacceptable impact under section 4.15 (79C(b)) of the Environmental Planning and Assessment Act 1979 or a significant impact under Section 7.3 of the Biodiversity Conservation Act 2016.

The proposal will not have a significant impact to terrestrial biodiversity and meets the requirement of clause 6.5 of the MLEP.

We recommend that the ameliorative conditions and management recommendations in this report be followed to limit disturbance during construction and to further reduce the impact of the proposal on potential bandicoot habitat and access.

6/5/21 Page 30 of 36



8 Ameliorative Conditions

- Access be provided from Bower Street by a 120mm gap under the full length of the base of the re-configured front gate. This is a lower than usual height recommended because the owner has small dogs that may fit under a 150mm gap.
- Access to the rear yards of the property is to be maintained along the eastern side of the proposed dwelling.
- Any gates along the through access path from the front to rear are to have 120mm gaps for bandicoots.
- Where there is no change in soil levels, bandicoot access is to be maintained through the
 existing gaps in the fences along the northern and southern boundaries.
- Replacement boundary fences are to have 120mm-300mm gaps every two metres along the fence (see Photo Page 1 for examples).
- It is recommended that any new replacement internal gates have gaps of at least 120mm under the full length of the gate.
- If any Bandicoot enters the site during works, works must be stopped until the Bandicoot has safely exited the site.
- Any injured or dead Long-nosed Bandicoots within the site must be reported National Wildlife Services or Northern Beaches Council.
- If the plans change from what is described in this report or what is shown on Map 1 then the impacts will change, and this report may need reviewing.
- In areas of habitat for the long-nosed bandicoot, landscape design should include native
 plant species to provide new and/or improved low dense clumping habitat to provide for
 potential foraging and nesting. The planting schedule should comprise species such as
 Lomandra sp. Dianella sp., Banksia spinulosa, Caustis sp., Xanthorrhoea sp., Isolepis sp.,
 Juncus sp., Calochlaena sp., Callistemon sp., Gleichenia sp. and Grevillea 'Robyn Gordon'
 (Manly DCP 2013 Section 3.3.1 [a][iv]).
- To the untrained eye an Endangered Long-nosed Bandicoot may be mistaken for a rat. To avoid direct physical harm to Long-nosed Bandicoots, it is important that workers on the site are aware of their presence and their conservation significance and the steps to take to protect them.
- While temporary fencing around the construction area is usually a standard requirement, even purpose-built fencing has been known to be ineffective in excluding bandicoots from construction sites. It is therefore essential that daily checks be undertaken to ensure the construction/works areas are free of bandicoot occupation including sheltering sites. All excavations and stockpiles of construction material are to be inspected daily prior to commencing operation to ensure that no bandicoots are sheltering in these areas. In the case that a Long-nosed Bandicoot is encountered within one of these work areas, no work shall proceed until the bandicoot has safely vacated the works area.
- Noise and vibration discourage bandicoot occupation of this and adjacent sites. Normal
 construction hours are to be adhered to, with no machinery to be used outside the hours of
 7:30am and 4:30pm.
- Bright lighting discourages bandicoot occupation. No bright lighting or motion detector lights are to be installed to illuminate the lawn or garden areas. A modest amount of low lighting is acceptable for safety purposes only.

6/5/21 Page 31 of 36



9 Management Recommendations

- The Companion Animals Act 1998 requires that dog and cat owners ensure that their cat or dog does not threaten or harm a person or animals.
- Domestic cats and dogs should be kept indoors from dusk to dawn. Cats should only be allowed outside if in a sealed cat run. Example solutions can be seen at these internet addresses:

http://www.catnip.com.au/design ideas.html

http://www.catnets.com.au/

http://www.catcagesaustralia.com.au/gallery/index.html

http://www.cat-world.com.au/cat-worldenclosures.htm

http://www.catmax.com.au/photo-gallery.php

- Dog owners must take all reasonable precautions to prevent your dog from escaping from the property on which it is being kept. If you fail to comply with this requirement, you may be liable for a penalty of \$880.
- Cats should be kept indoors at night, as there are benefits to both the cat and the community. Yowling and fighting is more of a problem at night. The noise is likely to be intrusive and may keep your neighbours awake. Keeping your cat indoors at night is recommended in the interests of both your cat's safety and community harmony. Many kinds of native wildlife are more active or more vulnerable to hunting at night. There is also evidence that cats hunt more during the night. By keeping your cat indoors, you can help reduce the number of native birds and animals that are killed in your area. Kittens can quickly become accustomed to staying indoors at night. Consider also containing your cat in a cat enclosure on your premises both during the day and during the night. Council Rangers can issue nuisance orders to cat owners.
- Injured bandicoots should be given expert care in order that they can be rehabilitated and returned to the population where possible. It is also important that any deceased bandicoots are reported, so that appropriate investigations can be undertaken to understand the cause of death to inform the future management and recovery of the endangered Long-nosed bandicoot population. Any injured or dead Long-nosed Bandicoots should be reported by phoning Council on 9976 1500 or Office of the Environment and Heritage (OEH) on 131 555
- Modest, low external lighting in the garden should be used at the minimum level required for safety.
- Landscaping watering and additional cover in the form of planting low, dense vegetation will
 increase the value of the foraging habitat and facilitate bandicoot persistence in the urban
 mosaic.
- Bandicoots and other native animals should not be fed artificial foods as it may cause them nutritional problems and may increase predation.
- Feral animals including cats and/or foxes should never be fed, nor should food be left out
 where they can access it, such as rubbish bins without lids, or in pet food bowls, as these
 animals present a significant threat to Long-nosed Bandicoots and other wildlife.
- Rat baiting is to only occur within buildings. No Rat baiting should occur under or around houses.
- Temporary chain wire fencing (minimum 300mm in height is sufficient) is to be installed around all work areas prior to commencement of works. Fencing is to be maintained for the

GIS Environmental Consultants

6/5/21 Page 32 of 36

- duration of the works. The fence is to be designed to minimise the possibility if Long-nosed Bandicoots accessing works areas.
- The use of insecticides, fertilisers, or snail baits should be avoided on the property. Garden insects will be kept in low numbers if Long-nosed Bandicoots are present.
- When the final North Head Long-nosed Bandicoot Recovery Plan is released it should be implemented where relevant.
- Care should be taken when driving in the area, especially at night as bandicoots have little road sense and cars are a major threat to bandicoots.
- Dead bandicoots should be reported by phoning Council on 9976 1500 as they will assist in monitoring the program.
- Please report all sightings of feral rabbits, feral or stray cats and/or foxes to Council on 9976 1500 or NPWS (OEH) on 9997 6102.



6/5/21 Page 33 of 36

Photo Page 3. Examples of Gaps in Boundary Fence



Figure 1. Example of gap under boundary fence for bandicoot access



Figure 2. Example of gap under boundary fence for bandicoot access



Figure 3. Example of gap under boundary fence for bandicoot access

GIS Environmental Consultants

6/5/21 Page 34 of 36

10 References and Relevant Literature

Banks, P. B. (2000). Population viability analysis for the Long-nosed Bandicoot population at North Head, NSW: modelling the effects of increased traffic flow on adult mortality. School of Biological Sciences, University of Sydney.

Banks, P. B. (2004). Population viability analysis in urban wildlife management: modelling management options for Sydney's quarantined bandicoots. School of Biological, Earth and Environmental Sciences, University of New South Wales.

Chambers, L. K. and Dickman, C. R. (2002) Habitat selection of the long-nosed Bandicoot *Perameles nasuta* (Mammalia, Peramelidae), in a patchy urban environment. *Austral Ecology* 27: 334-342.

Claridge, A. W. (1993). Fungal diet of the Long-nosed Bandicoot (*Perameles nasuta*) in south-eastern Australia. *Victorian Naturalist* 110: 86-91.

Cumberland Ecology (2016) Long-nosed Bandicoot Urban Monitoring Program 2016 - Interim Report

Department of Environment, Climate Change and Water (DECCW), Threatened Species Assessment Guidelines for the Assessment of Significance, dated August 2007

Office of Environment and Heritage (OEH), Threatened Species database, http://www.threatenedspecies.environment.nsw.gov.au/tsprofile/index.aspx

Hope B. 2012. Short-term response of the Long-nosed Bandicoot (*Perameles nasuta*) and the Southern Brown Bandicoot (*Isoondon obesulus*), to low-density prescribed fire in heathland vegetation, Wildlife Research 39:731-744.

Hughes, N. K. and Banks, P. B. (2006) An analysis of the May 2006 Census of the North Head Long-nosed Bandicoot Population: A report for DEC Central Directorate threatened Species Unit. School of Biological, Earth and Environmental Sciences, University of New South Wales.

Hughes, N. K. and Banks, P. B. (2010) Heading for greener pastures? Defining the foraging preferences of urban long-nosed bandicoots. School of Biological, Earth and Environmental Sciences, University of New South Wales.

Hughes, N. K. and Banks, P. B. (2013) An analysis of the 2012/2013 Census of the North Head Long-nosed Bandicoot Population: A report for OEH Central Directorate threatened Species Unit. School of Biological, Earth and Environmental Sciences, University of New South Wales.

Lenehan, J. and Banks, P. B. (2004). An analysis of the May 2004 North Head Long-nosed Bandicoot Population Census: A report for the NPWS Central Directorate Threatened Species Unit. School of Biological, Earth and Environmental Sciences, University of Sydney.

Menkhorst, P. W. & Knight, F. A. (2004). Field guide to the mammals of Australia. Oxford University Press, Melbourne.

NPWS. (2002). Endangered Population of Long-nosed Bandicoots (Perameles nasuta) at North Head, Draft Recovery Plan. NSW NPWS Hurstville.

NSW Scientific Committee (last amended June 2003). Final determination for the North Head population of the Long-nosed Bandicoot, Perameles nasuta.

Scott, L. K. (1995). Nutritional ecology and population biology of the Long-nosed Bandicoot (Perameles nasuta): Implications for conservation. School of Biological Sciences, University of Sydney.

Scott, L. K., Hume, I. D. & Dickman, C. R. (1999). Ecology and population biology of long-nosed bandicoots (Perameles nasuta) at North Head, Sydney National Park. Wildlife Research 26: 805-821.

Skelton, N., O. Richmond, A. Gilson and P. Wong, 2003, Fauna of North Head, GIS Environmental Consultants, North Curl Curl.

Skelton, N., P. Wong and E. Donner, 2004, Fauna and Fauna of Manly Councils Bushland Reserves, GIS Environmental Consultants, North Curl Curl.

Swimming Pools Act 1992, regulation 2018 and AS196.1-2012

Stoddart, E. (1995) "Long-nosed Bandicoot", pp. 184-185 in Mammals of Australia (Ed. R. Strahan). Reed Books, Chatswood.



6/5/21 Page 35 of 36





6/5/21 Page 36 of 36